

Section 3.4

Assessing barriers and facilitators to knowledge use

France Légaré, MD, PhD

Canada Research Chair in Implementation of Shared
Decision Making in Primary Care

Centre de recherche, Hôpital St-François d'Assise

Department of Family Medicine and Emergency Medicine,
Université Laval, Québec, Canada



Overview of chapter

- Why is it relevant to address barriers and facilitators to knowledge use?
- What are the key concepts and conceptual models for assessing barriers and facilitators to knowledge use?
- What are some tools for assessing barriers and facilitators to knowledge use?
- Future research



Key learning points

- Barriers and facilitators are the salient beliefs of self-efficacy, the most important determinant of behaviour change after intention.
- Taxonomies for barriers and facilitators have been developed and should be used when developing a knowledge-to-action project.
- Existing taxonomies should be further evaluated in other settings and contexts.
- Existing scales to assess barriers and facilitators to knowledge use need to be further evaluated.




Why is it relevant?

Diagnostic analysis:
N=329 physicians
Self-reported barriers explained 39% of the self-reported performance.

IMPROVING PATIENT CARE

The Implementation of Change in Clinical Practice



In the health care, a very large number of valuable insights, procedures and technologies become available each year. These come from well planned scientific research or careful experiments and evaluation in everyday practice. As many of these innovations are not adopted into daily practice, at the expense of improved patient care, it is crucial to understand how successful implementation can be achieved.

This book examines the process of introducing change and innovation into practice. With chapters on identifying problems in the implementation, selecting strategies, effective implementation of policies, and evaluation of the success of the changes, this book provides a



Why is it relevant?

- 78 studies of behaviour change in healthcare providers (Godin et al. 2008):
 - 72: determinants of intention
 - 16: determinants of behaviour.
- Factors most consistently associated with prediction of healthcare provider's behaviours and intention (*i.e.*, at least 50% of the time)
 - intention
 - **beliefs about capabilities (perceived behavioural control)**



KEY CONCEPTS AND CONCEPTUAL MODELS FOR ASSESSING BARRIERS AND FACILITATORS TO KNOWLEDGE USE





Clinical Practice Guidelines Framework for Improvement

Cabana et al. 1999

- This framework was based on an extensive search of the literature of barriers to physician adherence to CPGs and was organized according to **knowledge**, **attitudes**, or physician **behavior**.
- 76 published studies describing at least one barrier to adherence to CPGs.
- 293 potential barriers to physician guideline adherence.



Barriers and Facilitators to the implementation of shared decision making

Légaré et al. 2006

- Primary health care professional's views on barriers and facilitators to the implementation of the Ottawa Decision Support Framework in practice
- Extension of Cabana's model:
 - A specific definition was identified for each type of barrier;
 - Inclusion of a list of potential facilitators of knowledge use in clinical practice.





Barriers and Facilitators to the implementation of shared decision making

Légaré et al. 2006

- Cabana's model also extended with the attributes of innovation as proposed by the **Diffusion of Innovation theory** (*Rogers, 1995*).





A systematic review of health professionals' perceptions


Légaré et al. 2008

- Application of the revised version of the Clinical Practice Guideline Framework for Improvement.
- Barriers and facilitators to implementing shared decision making in clinical practice.
- 41 publications covering 38 unique studies.



Barriers and facilitators to SDM

Stages of the review process



(Légaré, Ratté, Gravel, Graham. Pt Educ Counseling, 2008)


Health professionals surveyed

Professionals	Number of Participants	%
Physicians	3253	90
Other professionals	341	9
Others	39	1
Total	3633	100


n=38 studies

*5 studies did not provide this information

Barriers



Facilitators



(Légaré, Ratté, Gravel, Graham. Pt Educ Counseling, 2008)



TOOLS FOR ASSESSING BARRIERS AND FACILITATORS TO KNOWLEDGE USE



CIHR IRSC

Canada 

Tool for assessing barriers to adherence to hand hygiene guidelines

Larson, 2004

- Was developed and tested on a group of 21 infectious disease clinicians.
- 6-point Likert scale
- 2 sections:
 - Attitudinal statements about practice guidelines in general
 - Specific statements regarding the Hand Hygiene Guideline





Instrument to assess barriers and facilitators to knowledge use

Wensing et Grol, 2005

- Applied to 12 different implementation studies in the Netherlands.
- Literature analyses and focus groups with implementation experts to identify possible barriers to change.
- Validation studies to test psychometric characteristics of the questionnaires.



BARRIERS Scale

Funk et al. 1991

- Develop to assess barriers to research utilization based on four key dimensions: nurse, setting, research and presentation.
- 29 items
- 4 subscales that map 4 key dimensions :
 - (1) characteristics of the adopter;
 - (2) characteristics of the organization;
 - (3) characteristics of the innovation; and
 - (4) characteristics of the communication





Knowledge
to action

Des connaissances
à la pratique

FUTURE RESEARCH



CIHR IRSC

Canada 

Challenges that will need to be addressed

- Need to standardize the reporting of barriers and facilitators to translating research into clinical practice.
- Need to address barriers as well as facilitators to knowledge use because one factor can be perceived as both a barrier and a facilitator.
- Need to adapt existing instruments to the assessment of facilitators of knowledge.
- Need to test existing instruments in diverse clinical and cultural contexts.



Research questions

1. How should we measure barriers and facilitators to research use to infer what the collective is thinking?
2. Should we collect data from individuals to make sense of the group's thinking?
3. If we do, should we use the mean or the median to represent the group or should we focus on the variation or only on the outliers?
4. How many individuals in a group have to perceive something is a barrier before we decide to address it with an intervention?
5. Is the perception of the opinion leader the most important one?



References

1. Cabana M, Rand C, Powe N, Wu A, Wilson M, Abboud P, et al. Why don't physicians follow clinical practice guidelines? A framework for improvement. *JAMA*. 1999;282:1458-65.
2. Funk SG, Champagne MT, Wiese RA, Tornquist EM. BARRIERS: the barriers to research utilization scale. *Appl Nurs Res*. 1991 Feb;4:39-45.
3. Larson E. A tool to assess barriers to adherence to hand hygiene guideline. *American journal of infection control*. 2004 Feb;32:48-51.
4. Légaré F, O'Connor A M, Graham ID, Saucier D, Cote L, Blais J, et al. Primary health care professionals' views on barriers and facilitators to the implementation of the Ottawa Decision Support Framework in practice. *Patient Educ Couns*. 2006 Nov;63:380-90.
5. Légaré F, Ratté S, Gravel K, Graham ID, Barriers and facilitators to implementing shared decision making in clinical practice: update of a systematic review of health professionals perceptions. *Patient Educ Couns*. 2008; 73:526-35.
6. Rogers EM. Diffusion of innovations. Fourth ed. New York: The Free Press 1995.
7. Wensing M, Grol R. Methods to identify implementation problems. In: Grol R, Wensing M, Eccles M, eds. Improving patient care The implementation of change in clinical practice. Oxford: Elsevier Butterworth Heinemann 2005:109-21
8. Godin G, Bélanger-Gravel A, Eccles M, Grimshaw J. Healthcare professionals' intentions and behaviours: a systematic review of studies based on social cognitive theories. *Implement Sci*. 2008 Jul 16;3:36.

